

# The discrete Moser–Veselov algorithm for the free rigid body

Antonella Zanna and Robert McLachlan

Report in Informatics nr. 255 (2003). To appear in Found. Comp. Math.

Antonella Zanna  
University of Bergen, Norway

email: [anto@ii.uib.no](mailto:anto@ii.uib.no)  
<http://www.ii.uib.no/~anto>

Universitet i Bergen



1/100



Back

Close

Report Documentation Page				Form Approved OMB No. 0704-0188	
Public reporting burden for the collection of information is estimated to average 1 hour per response, including the time for reviewing instructions, searching existing data sources, gathering and maintaining the data needed, and completing and reviewing the collection of information. Send comments regarding this burden estimate or any other aspect of this collection of information, including suggestions for reducing this burden, to Washington Headquarters Services, Directorate for Information Operations and Reports, 1215 Jefferson Davis Highway, Suite 1204, Arlington VA 22202-4302. Respondents should be aware that notwithstanding any other provision of law, no person shall be subject to a penalty for failing to comply with a collection of information if it does not display a currently valid OMB control number.					
1. REPORT DATE <b>03 JAN 2005</b>		2. REPORT TYPE <b>N/A</b>		3. DATES COVERED <b>-</b>	
4. TITLE AND SUBTITLE <b>The discrete MoserVeselov algorithm for the free rigid body</b>				5a. CONTRACT NUMBER	
				5b. GRANT NUMBER	
				5c. PROGRAM ELEMENT NUMBER	
6. AUTHOR(S)				5d. PROJECT NUMBER	
				5e. TASK NUMBER	
				5f. WORK UNIT NUMBER	
7. PERFORMING ORGANIZATION NAME(S) AND ADDRESS(ES) <b>University of Bergen, Norway</b>				8. PERFORMING ORGANIZATION REPORT NUMBER	
9. SPONSORING/MONITORING AGENCY NAME(S) AND ADDRESS(ES)				10. SPONSOR/MONITOR'S ACRONYM(S)	
				11. SPONSOR/MONITOR'S REPORT NUMBER(S)	
12. DISTRIBUTION/AVAILABILITY STATEMENT <b>Approved for public release, distribution unlimited</b>					
13. SUPPLEMENTARY NOTES <b>See also ADM001749, Lie Group Methods And Control Theory Workshop Held on 28 June 2004 - 1 July 2004., The original document contains color images.</b>					
14. ABSTRACT					
15. SUBJECT TERMS					
16. SECURITY CLASSIFICATION OF:			17. LIMITATION OF ABSTRACT <b>UU</b>	18. NUMBER OF PAGES <b>31</b>	19a. NAME OF RESPONSIBLE PERSON
a. REPORT <b>unclassified</b>	b. ABSTRACT <b>unclassified</b>	c. THIS PAGE <b>unclassified</b>			



# Overview

The subject of this talk is the numerical solution of the free RB equations,

$$M' = [M, \Omega], \quad M = \Omega J + J \Omega,$$

where  $M, \Omega$  are skew-symmetric matrices and  $J$  is a diagonal matrix with positive entries.

$M$  is the matrix of **body momenta**

$\Omega$  is the matrix of **body angular velocity**

Often the above equations are associated with the equations that give the configuration of the body in the fixed frame,

$$Q' = Q\Omega, \quad Q \in \text{SO}(N).$$

- The Discrete Moser–Veselov description of the rigid body
- Integrability of the discrete algorithm
- Backward error analysis of the the DMV algorithm
- Higher order integrable approximations
- Numerical experiments and comparisons with other methods
- Explicit methods for the  $3 \times 3$  case



Back

Close

# The Moser–Veselov discrete version of the dynamics of a Rigid Body

The Lagrangian of the continuous RB equations, is the kinetic energy,

$$L = \frac{1}{2} \text{tr}(\Omega^\top M) = \frac{1}{2} \text{tr}(-\Omega^2 J - \Omega J \Omega) = \text{tr}(\Omega^\top J \Omega), \quad (1)$$

where we take into account that  $\Omega^\top = -\Omega$  and that the trace is invariant under cyclic permutations. Following (Marsden, Pekarsky & Shkoller 1999), discretise  $\Omega = X^{-1} \dot{X}$ , where  $X \in \text{SO}(N)$  is the configuration of the body, using a finite difference approximation of the derivative,

$$\Omega = X^{-1} \dot{X} \approx \frac{1}{h} X_{k+1}^\top (X_{k+1} - X_k), \quad X_k, X_{k+1} \in \text{SO}(N),$$

which gives

$$L \approx \frac{1}{h^2} \text{tr}(J - X_k^\top X_{k+1} J - J X_{k+1}^\top X_k - X_k^\top X_{k+1} J X_{k+1}^\top X_k).$$

Due to the orthogonality of the  $X_k$ 's and the cyclicity of the trace, the first and the last term cancel, and moreover, we can write

$$L \approx \frac{2}{h^2} \text{tr}(X_k J X_{k+1}^\top).$$

Up a scaling factor, this is precisely the discrete Lagrangian of Moser and Veselov.



Consider the functional  $S(X)$  determined by

$$S = \sum_k \text{tr}(X_k J X_{k+1}^\top)$$

where  $X = \{X_k\}$  with  $X_k \in O(N)$  and  $J$  is a diagonal matrix. To obtain the stationary points of  $S$ , we consider

$$\sum_k \text{tr}(X_k J X_{k+1}^\top) - \frac{1}{2} \sum_k \text{tr}(\Lambda_k (X_k X_k^\top - I)),$$

(where  $\Lambda_k = \Lambda_k^\top$  is a Lagrange multiplier), and  $\delta S = 0$  becomes

$$X_{k+1} J + X_{k-1} J = \Lambda_k X_k,$$

from which, multiplying by  $X_k^\top$  on the right and taking into consideration the symmetry of  $\Lambda_k$ ,

$$X_{k+1} J X_k^\top + X_{k-1} J X_k^\top = \Lambda_k = \Lambda_k^\top = X_k J X_{k+1}^\top + X_k J X_{k-1}^\top, \quad (2)$$

hence, the *discrete analogue of the angular momentum in space*,

$$m_k = X_k J X_{k-1}^\top - X_{k-1} J X_k^\top,$$

**is conserved.**





In the body variables, setting  $\omega_k = X_k^\top X_{k-1} \in O(N)$  and  $M_k = X_{k-1}^{-1} m_k X_{k-1} = \omega_k^\top J - J \omega_k \in \mathfrak{so}(N)^*$  (angular momentum w.r.t. the body), (2) becomes

$$\begin{aligned} M_{k+1} &= \omega_k M_k \omega_k^\top \\ M_k &= \omega_k^\top J - J \omega_k. \end{aligned} \tag{3}$$

the **discrete Euler–Arnold** equation.

In the continuous limit: when  $t_k = t_0 + k\varepsilon$ ,  $k = 0, 1, 2, \dots$ ,

- $X_k = X(t_k)$
- $\omega_k = X_k^\top X_{k-1} \approx I - \varepsilon \Omega(t_k)$ ,
- $M_k \approx \varepsilon(J\Omega + \Omega J) = \varepsilon M(t_k)$ ,

letting  $\varepsilon \rightarrow 0$ , one obtains the familiar Euler–Arnold equations for the motions of the  $N$ -dimensional rigid body,

$$\begin{aligned} M' &= [M, \Omega] \\ M &= J\Omega + \Omega J, \quad \Omega \in \mathfrak{so}(N). \end{aligned}$$



To solve the discrete Euler–Arnold equations (3):

- For  $k = 0, 1, 2, \dots$ , find  $\omega_k \in \text{SO}(N)$  such that  $M_k = \omega_k^\top J - J \omega_k$ .
- Update  $M_{k+1} = \omega_k M_k \omega_k^\top$ .

By construction, this algorithm

- preserves exactly momentum and energy (integrable map)
- is a second order approximation to the continuous rigid body
- preserves the standard Poisson structure of  $T^*\mathfrak{so}(N)$ ,

$$\{f, g\} = \text{tr}(M[f_M, g_M]), \quad f, g \in C^\infty(\mathfrak{so}(N)),$$

where  $f_M = (\partial f / \partial M_{i,j})$ .



Note that

- Also the IMR is second order, preserves all the integrals of the continuous rigid body.
- Another much used method is a Lie–Poisson integrator of McLachlan and Reich (LP2). For the  $3 \times 3$  RB, it consists in splitting the Hamiltonian

$$H = \mathcal{H}_1 + \mathcal{H}_2 + \mathcal{H}_3 = \frac{m_1^2}{J_2 + J_3} + \frac{m_2^2}{J_1 + J_3} + \frac{m_3^2}{J_1 + J_2}$$

and integrating explicitly (a la Strang) the vector fields of each split Hamiltonian. The method is second order, explicit, preserves the Poisson structure but does not preserve  $H$ .





Error in the Hamiltonian function  $H$  in the interval  $[0, 100]$  and for  $h = \frac{1}{2}$ .

The components of the vector  $\mathbf{m}_k$  for  $h = \frac{1}{2} \dots$

$$\mathbf{m} = \begin{bmatrix} m_1 \\ m_2 \\ m_3 \end{bmatrix} \equiv \hat{\mathbf{m}} = M = \begin{bmatrix} 0 & -m_3 & m_2 \\ m_3 & 0 & -m_1 \\ -m_2 & m_1 & 0 \end{bmatrix}$$







# Integrability of the Moser–Veselov equation

Recall the Moser–Veselov equation,

$$M_k = \omega_k^\top J - J \omega_k, \quad M_k^\top = -M_k, \quad \omega_k^\top \omega_k = I. \quad (4)$$

in tandem with the update  $M_{k+1} = \omega_k M_k \omega_k^\top$ .

- The Moser–Veselov equation (4) has not a unique solution;

**Lemma 1** (*Moser and Veselov*) Equation (4) is equivalent to the factorization

$$(I - \lambda M_k - \lambda^2 J^2) = (\omega_k^\top + \lambda J)(\omega_k - \lambda J)$$

Rewrite the above factorization as

$$(I - \lambda M_k - \lambda^2 J^2) = A_k^\top(\lambda) A_k(\lambda) \quad A_k(\lambda) = (\omega_k - \lambda J)$$

and define the mapping with image point  $M_{k+1}$  such that

$$(I - \lambda M_{k+1} - \lambda^2 J^2) = A_k(\lambda) A_k^\top(\lambda).$$

By construction,

$$(I - \lambda M_{k+1} - \lambda^2 J^2) = A_k(\lambda)(I - \lambda M_k - \lambda^2 J^2)A_k^{-1}(\lambda),$$

hence the map is an **integrable** as it is an isospectral flow.





# BEA for DMV

Recall the DMV equations and the continuous RB equations

$$\begin{aligned} M_{k+1} &= \omega_k M_k \omega_k^\top, & M' &= [M, \Omega] \\ M_k &= \omega_k^\top J - J \omega_k, & M &= \Omega J + J \Omega, \end{aligned}$$

where  $\omega_k \approx I - h\Omega(t_k)$ .

We wish to write

$$M_{k+1} = \Phi_h(M_k) = M_k + h[M_k, \Omega_k] + h^2 d_2 + h^3 d_3 + h^4 d_4 + \dots,$$

and find the modified vector field

$$\tilde{M}' = [\tilde{M}, \tilde{\Omega}] + h f_2(\tilde{M}, \tilde{\Omega}) + h^2 f_3(\tilde{M}, \tilde{\Omega}) + h^3 f_4(\tilde{M}, \tilde{\Omega}) + \dots \quad (5)$$

such that  $\Phi_h(M_k)$  equals the solution  $\tilde{M}(t_{k+1})$  at time  $t_{k+1} = t_0 + (k+1)h$  of the modified vector field (5).

To find  $\Phi_k(h)$ , we write

$$\omega_k = \exp(-h\Omega_0 - h^2\Omega_1 - h^3\Omega_2 - h^4\Omega_3 - h^5\Omega_4 + \dots), \quad (6)$$

where  $\Omega_0, \Omega_1, \Omega_2, \dots$ , are skew-symmetric matrices computed so that

$$\omega_k^\top J - J \omega_k = h(\Omega(t_k)J + J\Omega(t_k)). \quad (7)$$



we obtain

$$\begin{aligned} h(\Omega(t_k)J + J\Omega(t_k)) &= h(\Omega_0J + J\Omega_0) + h^2(\Omega_1J + J\Omega_1 + \tfrac{1}{2}(\Omega_0^2J - J\Omega_0^2)) \\ &\quad + h^3(\Omega_2J + J\Omega_2 + \tfrac{1}{2}[(\Omega_0\Omega_1 + \Omega_1\Omega_0), J] + \tfrac{1}{6}(\Omega_0^3J + J\Omega_0^3)) + \dots \end{aligned}$$

Comparing left and right-hand-sides, it is trivially observed that the order- $h$  term disappears if  $\Omega_0 = \Omega$  (to simplify notation, we omit the dependence of  $\Omega$  on  $t_k$ ). In order to annihilate the  $h^2$ -term, we require that

$$\Omega_1J + J\Omega_1 + \frac{1}{2}(\Omega_0^2J - J\Omega_0^2) = 0.$$

Recall that  $M = \Omega J + J\Omega$  and hence  $M' = \Omega'J + J\Omega'$ . On the other hand,  $M' = [M, \Omega] = -(\Omega^2J - J\Omega^2)$ . Hence we can write

$$O = \Omega_1J + J\Omega_1 - \frac{1}{2}M' = \Omega_1J + J\Omega_1 - \frac{1}{2}(\Omega'J + J\Omega')$$

and the identity is satisfied by if and only if

$$\Omega_1 = \frac{1}{2}\Omega'. \quad (8)$$

In general, the **algorithm** to derive  $\Omega_i$ , for  $i = 1, 2, \dots$ , is

1. Find the coefficient of  $h^{i+1}$  in (7) and set it equal to zero. This will give an equation of the type  $\Omega_iJ + J\Omega_i = C_iJ + JC_i + [D_i, J]$ . Note that the terms  $C_iJ + JC_i$  have an odd occurrence of the  $\Omega_j$ s, while the terms of the type  $[D_i, J]$  have an even occurrence of the  $\Omega_j$ s.
2. Use the derivatives of  $M$  and  $\Omega$  to express the term  $[D_i, J]$  as  $\tilde{C}_iJ + J\tilde{C}_i$ .



3. Deduce  $\Omega_i = C_i + \tilde{C}_i$ .

---


$$\begin{aligned}\Omega_0 &= \Omega \\ \Omega_1 &= \frac{1}{2}\Omega' \\ \Omega_2 &= \frac{1}{6}\Omega'' - \frac{1}{6}\Omega^3 \\ \Omega_3 &= \frac{1}{8}\Omega''' - \frac{1}{24}(5\Omega^2\Omega' + 2\Omega\Omega'\Omega + 5\Omega'\Omega^2)\end{aligned}$$


---

The functions  $\Omega_i$

Once the  $\Omega_i$ s are known, substituting back in  $M_{k+1} = \omega_k^\top M_k \omega_k$  and using the well known identity

$$\exp(X)Y \exp(-X) = \exp_{\text{ad}_X} Y = \sum_{k=0}^{\infty} \frac{1}{k!} \text{ad}_X^k(Y),$$

where  $\text{ad}_X(Y) = [X, Y]$  and, recursively,  $\text{ad}_X^k(Y) = [X, \text{ad}_X^{k-1}(Y)]$ , we find the expressions for the functions  $d_i$  in terms of the  $\Omega_{i-1}, \Omega_{i-2}, \dots, \Omega_0$ ,

$$d_i = \sum_{j=1}^i \frac{(-1)^j}{j!} \sum_{k_1+k_2+\dots+k_j=i-j} \text{ad}_{\Omega_{k_1}} \text{ad}_{\Omega_{k_2}} \cdots \text{ad}_{\Omega_{k_j}} M, \quad k_1, \dots, k_j \in \{0, 1, \dots, i-1\}. \quad (9)$$

$$\begin{aligned}d_2 &= \frac{1}{2}([M, \Omega'] + [[M, \Omega], \Omega]), \\ d_3 &= \frac{1}{4}[M, \Omega''] + \frac{1}{4}[[M, \Omega'], \Omega] + \frac{1}{4}[[M, \Omega], \Omega'] + \frac{1}{6}[[[M, \Omega], \Omega], \Omega] - \frac{1}{6}[M, \Omega^3], \\ d_4 &= \dots,\end{aligned} \quad (10)$$



## Taylor expansion of the solution of the modified equation

Consider

$$\frac{d}{dt}\tilde{y} = f(\tilde{y}) + hf_2(\tilde{y}) + h^2f_3(\tilde{y}) + \dots,$$

where  $f(M) = [M, \Omega] = [M, \mathcal{J}^{-1}M]$  is the original vector field of the RB equations, where  $\mathcal{J}$  is a linear operator, defined such that  $\mathcal{J}\Omega = \Omega\mathcal{J} + \mathcal{J}\Omega = M$ . Putting  $\tilde{y}(t) = M(t)$ , we expand the solution of the above equation in a Taylor series and collect corresponding powers of  $h$ ,

$$\begin{aligned}\tilde{y}(t+h) &= M(t) + hf(M) + h^2 \left( f_2(M) + \frac{1}{2!}f'f(M) \right) \\ &\quad + h^3 \left( f_3(M) + \frac{1}{2!}(f'f_2(M) + f'_2f(M)) + \frac{1}{3!}(f''(f, f)(M) + f'f'f(M)) \right) + \dots,\end{aligned}$$

where  $f'$  is considered as a linear operator,  $f''$  as a bilinear operator and so on and so forth. In our case,

$$\begin{aligned}f'(z)(M) &= [z, \mathcal{J}^{-1}M] + [M, \mathcal{J}^{-1}z] \\ &= [z, \Omega] + [M, \mathcal{J}^{-1}z] \\ f''(z_1, z_2)(M) &= 2[z_1, \mathcal{J}^{-1}z_2],\end{aligned}$$

and, since  $f$  is quadratic,  $f'''$  and all the other higher derivatives equal zero.

At this point it is important to stress an important difference between the expressions for the modified vector field of (Hairer, Lubich & Wanner 2002) and ours. While the vector field discussed in (Hairer et al. 2002) is in  $\mathbb{R}^n$ , hence the  $f''$  is a symmetric quadratic operator, this is not the case for our vector field which is on matrices, thus

$$f''(f'f, f) \neq f''(f, f'f).$$




This non-commutative case is discussed with more generality in (Munthe-Kaas & Krogstad 2002). However, we observe that *all* the terms containing combinations of  $f''$ ,  $f'$  and  $f$  correspond simply to higher derivatives of  $f$ . The mixed terms are treated instead specifically.

After some algebra, we have

$$\begin{aligned}
 f_2 &= d_2 - \frac{1}{2!} f' f(M) \\
 &= O, \\
 f_3 &= d_3 - \frac{1}{3!} (f''(f, f)(M) + f' f' f(M)) \\
 &= \frac{1}{12} [M, \Omega'' - [\Omega, \Omega'] - 2\Omega^3], \\
 f_4 &= d_4 - \frac{1}{4!} M^{(iv)} - \frac{1}{2!} (f' f_3 + f_3' f) \\
 &= O, \\
 f_5 &= d_5 - \frac{1}{5!} M^{(v)} - \frac{1}{2!} (f' f_4 + f_4 f' + \frac{1}{2!} \frac{d}{dt} (f_3' f + f' f_3)) \\
 &= \frac{1}{80} [M, \Omega^{(iv)}] - \frac{1}{80} [M, [\Omega, \Omega''']] + \frac{3}{40} [M, \Omega^5 - \Omega' \Omega \Omega'] \\
 &\quad + \frac{1}{80} [M, [\Omega', \Omega'']] - \frac{1}{40} [M, \Omega \Omega'' \Omega] - \frac{1}{20} [M, \Omega^2 \Omega'' + \Omega'' \Omega^2] \\
 &\quad + \frac{1}{20} [M, [\Omega^3, \Omega']] - \frac{1}{40} [M, \Omega \Omega'^2 \Omega + \Omega \Omega \Omega'^2 + \Omega [\Omega, \Omega'] \Omega].
 \end{aligned} \tag{11}$$





The DMV solution of the RB equations (dotted line), the exact solution (solid line) and the trajectories corresponding to the modified vector fields  $f + h^2 f_3$  (dashed line) and  $f + h^2 f_3 + h^4 f_5$  (dash-dotted line) in the interval  $[0, 50]$  with  $h = \frac{8}{10}$ .



Back

Close

## Some important results about DMV

**Theorem 2** *The DMV is time-reversible, hence  $f_{2i} = 0$ ,  $i = 1, 2, \dots$*

**Theorem 3** (Moser–Veselov) *In the  $3 \times 3$  case, the DMV is a time-reparametrisation of the flow of the original vector field of the rigid body.*

Since the mapping preserves the underlying Poisson structure and all the integrals  $F_i = c_i$  of the system, it commutes with all commuting Hamiltonian flows generated by the  $F_i$ s,  $M' = \{M, \nabla F_i\}$ . The nonsingular compact level sets  $T_c = \cap_i (F_i = c_i)$  consists of a finite union of 1-dimensional tori and on each torus the DMV mapping is a shift along the trajectory depending on the integral quantity  $H_2$ .

Hence, the DMV solves the modified equation

$$M' = (1 + h^2\tau_3 + h^4\tau_5 + \dots + h^{2i}\tau_{2i+1} + \dots)[M, \Omega],$$

where  $h$  is the stepsize of integration and the  $\tau_{2i+1}$ , for  $i = 1, 2, \dots$ , are constants.



15/100



Back

Close



Introduce the constants

$$\begin{aligned}C_{J,i,j} &= J_1^i J_2^j + J_1^i J_3^j + J_2^i J_3^j, \\C_{J,i} &= C_{J,i,i} \\C_J &= C_{J,1} \\\Delta &= (J_1 + J_2)(J_1 + J_3)(J_2 + J_3) \\H_2 &= (J_1 + J_2)(J_1 + J_3)(J_2 + J_3)H - C_J \|\mathbf{m}\|_2.\end{aligned}$$

**Theorem 4** *One has*

$$\tau_3 = \frac{1}{6\Delta^2}((3\det(J)\mathrm{tr}(J) + C_{J,2})\|\mathbf{m}\|_2^2 + (3C_J + \mathrm{tr}(J^2))H_2),$$

and

$$\begin{aligned}\tau_5 &= \frac{1}{40\Delta^4} \Big( (3\mathrm{tr}(J^4) + 27C_{J,2} + 15\mathrm{tr}(J^2)C_J + 45\det(J)\mathrm{tr}(J))H_2^2 \\&\quad + (10C_{J,3} + 50\det(J)\mathrm{tr}(J)C_J + 10\det(J)\mathrm{tr}(J)\mathrm{tr}(J^2) + 2C_{J,2}\mathrm{tr}(J^2) - 28\det(J^2))\|\mathbf{m}\|_2^2 H_2 \\&\quad + (60\det(J^2)C_J + 3C_{J,4} + 27\det(J^2)\mathrm{tr}(J^2) + 15\det(J)(C_{J,2,3} + C_{J,3,2}))\|\mathbf{m}\|_2^4 \Big).\end{aligned}$$

*Proof.* By direct computation of  $f_3$  and  $f_5$ . □



16/100

Universitet i Bergen



Back

Close



# Higher-order integrable methods

For the original RB equations, scaling the initial condition is equivalent to scaling time.

In our case, we know that DMV is a time-rescaling of the original RB equation. Therefore we wish to rescale the initial condition to obtain a better approximation of the unscaled original RB.

I.C. DMV

New I.C. DMV

$$h(\Omega(t_k)J + J\Omega(t_k)) \quad \frac{h(\Omega(t_k)J + J\Omega(t_k))}{1 + \tilde{\tau}_3 h^2 + \tilde{\tau}_5 h^4 + \dots}$$

We perform again the backward error analysis. We set now  $\tilde{\omega} = \exp(-h\tilde{\Omega}_0 - h^2\tilde{\Omega}_1 + \dots)$  and solve for the  $\tilde{\Omega}_i$ s as the skew-symmetric matrices that solve

$$h(1 - \tilde{\tau}_3 h^2 + (\tilde{\tau}_3^2 - \tilde{\tau}_5)h^4 + \dots)(\Omega J + J\Omega) = \tilde{\omega}^\top J - J\tilde{\omega}. \quad (12)$$

$$\begin{aligned} \tilde{f}_3 &= \tilde{d}_3 - \frac{1}{3!}M''' = -\tilde{\tau}_3[M, \Omega] + d_3 - \frac{1}{3!}M''' \\ &= -\tilde{\tau}_3[M, \Omega] + f_3 = (-\tilde{\tau}_3 + \tau_3)[M, \Omega], \end{aligned}$$

hence, in order to have an order-four scheme, we must set  $\tilde{f}_3 = 0$  which corresponds to the choice

$$\tilde{\tau}_3 = \tau_3.$$



Back

Close

After further computations, one has

$$\tilde{f}_5 = 0 \leftrightarrow \tilde{\tau}_5 = \tau_5 - 2\tau_3^2.$$

This value of  $\tilde{\tau}_5$  gives indeed a method of order six.

The new proposed algorithms of order four and six are described below.

### The DMV4 algorithm:

1. Compute  $\tau_3$  and set  $M_0 := M(t_0)h/(1 + h^2\tau_3)$ .
2. For  $k = 0, 1, \dots, n - 1$ ,  
    find the unique  $w_k$  as above such that  $M_k = \omega_k^\top J - J\omega_k$   
    set  $M_{k+1} = \omega_k M_k \omega_k^\top$   
    end
3. Reconstruct  $M_n := M_n(1 + h^2\tau_3)/h \approx M(t_n)$ .

### The DMV6 algorithm:

1. Compute  $\tau_3, \tau_5$  and set  $\tilde{\tau}_5 = \tau_5 - 2\tau_3^2$  and  $M_0 := M(t_0)h/(1 + h^2\tau_3 + h^4\tilde{\tau}_5)$ .
2. For  $k = 0, 1, \dots, n - 1$ ,  
    find the unique  $w_k$  as above such that  $M_k = \omega_k^\top J - J\omega_k$   
    set  $M_{k+1} = \omega_k M_k \omega_k^\top$   
    end
3. Reconstruct  $M_n := M_n(1 + h^2\tau_3 + h^4\tilde{\tau}_5)/h \approx M(t_n)$ .



## Some numerical experiments

We consider with initial condition

$$\mathbf{m}_0 = \begin{bmatrix} 0.4165 \\ 0.9072 \\ 0.0588 \end{bmatrix}$$

and matrix  $J$  given as

$$J = \begin{bmatrix} 0.9218 & 0 & 0 \\ 0 & 0.7382 & 0 \\ 0 & 0 & 0.1763 \end{bmatrix}$$

and compare the DMV explicit scheme with the Hamiltonian-splitting method LP2 of (McLachlan 1993)

$$H = \frac{m_1^2}{J_2 + J_3} + \frac{m_2^2}{J_1 + J_3} + \frac{m_3^2}{J_1 + J_2} = \mathcal{H}_1 + \mathcal{H}_2 + \mathcal{H}_3$$

and the Implicit Midpoint Rule (IMR),

$$\mathbf{m}_{k+1} = \mathbf{m}_k + hf\left(\frac{\mathbf{m}_k + \mathbf{m}_{k+1}}{2}\right),$$

where

$$\mathbf{f}(\mathbf{m}) = \mathbf{m} \times (\tilde{J})^{-1} \mathbf{m}, \quad \tilde{J} = \begin{bmatrix} J_2 + J_3 & 0 & 0 \\ 0 & J_1 + J_3 & 0 \\ 0 & 0 & J_1 + J_2 \end{bmatrix}.$$

Universitet i Bergen




19/100




Back

Close



Error versus step size computed at  $T = 100$  for the methods LP2, IMR, DMV, DMV4, DMV6.






Floating point operations versus accuracy ( $T = 100$ ) for the methods LP2, IMR, DMV, DMV4, DMV6. The roots of  $P(\lambda)$  are recomputed at each step, use QR with pivoting, (DMV  $\approx 22$  LP2 per step).



Back

Close



Floating point operations versus accuracy ( $T = 100$ ) for the methods LP2, IMR, DMV, DMV4, DMV6 and RATTLE6. The roots of  $P(\lambda)$  are computed once, use LU instead of QR (DMV  $\approx 19$ )



Back

Close

LP2 per step).

Method	$h = \frac{1}{16}$	$h = \frac{1}{2}$	$h = 1.2$	$h = 2.2$	$h = 2.5$	$h = 4$
LP2	6.7903e-03	5.1043e-01	1.6055e+00	1.7002e+00	1.9489e+00	1.3902e+00
IMR	1.5494e-04	9.9329e-03	1.3119e-01	3.9514e-01	2.5276e-01	6.1905e-01
DMV	1.5014e-02	5.9899e-01	NaN	NaN	NaN	NaN
DMV4	1.757e-07	7.6167e-04	1.0785e-01	1.6094e+00	5.0245e-01	5.1624e-01
DMV6	1.962e-10	1.6440e-06	7.0269e-02	NaN	7.0407e-01	1.0519e-01

Error for the various methods and selected step sizes



23/100

Universitet i Bergen



Back

Close



## Connections with matrix Riccati equations

Consider the matrix equation

$$M = XJ - JX^\top. \quad (13)$$

Cardoso & Leite (2003) shown that every solution of (13) (not necessarily orthogonal) is of the form

$$X = (M/2 + S)J^{-1},$$

for some symmetric matrix  $S$ .

Furthermore,  $X$  is a *orthogonal* solution of (13) if and only if  $S$  is a symmetric solution of the Riccati equation

$$S^2 + S(M/2) + (M/2)^\top S - (M^2/4 + J^2) = 0. \quad (14)$$

Riccati equations are associated to symplectic matrices. In our case, the symplectic matrix is

$$H_{\text{symp}} = \begin{bmatrix} \frac{M}{2} & I \\ \frac{M^2}{4} + J^2 & \frac{M}{2} \end{bmatrix}. \quad (15)$$

If  $\frac{M^2}{4} + J^2$  is positive definite, it has been shown in (Cardoso & Leite 2003) that (14) has a unique solution  $S$  which is symmetric, positive definite, and such that the eigenvalues of  $W = M/2 + S$  have positive real parts. This matrix  $W$  is precisely the same matrix in Moser & Veselov (1991), from which one obtains

$$\omega = WJ^{-1}.$$



**Algorithm**(Cardoso & Leite 2003): Compute  $X$ , the unique solution of (13) in the special orthogonal group  $SO(n)$ .

1. Find a real Schur form of  $H_{\text{symp1}}$ ,

$$\tilde{Q}^\top H_{\text{symp1}} \tilde{Q} = \begin{bmatrix} T_{11} & T_{12} \\ O & T_{22} \end{bmatrix}, \quad (16)$$

where  $T_{11}$  and  $T_{22}$  are block upper-triangular matrices such that the real parts of the spectrum of  $T_{11}$  are positive and the real parts of the spectrum of  $T_{22}$  are negative definite.

2. Partition  $\tilde{Q}$  accordingly,

$$\tilde{Q} = \begin{bmatrix} Q_{11} & Q_{12} \\ Q_{21} & Q_{22} \end{bmatrix}.$$

Then, compute

$$S = Q_{21} Q_{11}^{-1}.$$

3. Compute

$$X = \left( \frac{M}{2} + S \right) J^{-1}.$$

Some computational details

- Compute real Schur forms by QR iterations for eigenvalues (Golub & van Loan 1989)
- Cost:  $\mathcal{O}((2N)^3)$  operations (implicit methods for ODEs:  $\mathcal{O}(N^3)$ )

$N$  being the dimension of  $M$ .





## The case $N = 3$

In this case,

- it is possible to find an **explicit spectral decomposition** of  $H_{\text{symp1}}$  (without the QR eigenvalue method)
- construct the real Schur decomposition (16) and hence  $X$  from the eigenstructure of  $H_{\text{symp1}}$ .

This yields an **explicit** numerical method for the reduced RB equations.



The eigenvalues of the matrix  $H_{\text{sympI}}$ ,

$$H_{\text{sympI}} = \begin{bmatrix} \frac{M}{2} & I \\ \frac{M^2}{4} + J^2 & \frac{M}{2} \end{bmatrix} \quad (17)$$

are the solutions of the **quadratic eigenvalue problem**

$$P(\lambda) = \det(\lambda^2 I - \lambda M - J^2) = 0.$$

Without loss of generality, we assume that  $J$  is diagonal, with entries  $J_1, J_2, J_3$ . Then,

$$\begin{aligned} -P(\lambda) &= \lambda^6 - \lambda^4 (J_1^2 + J_2^2 + J_3^2 - m_{12}^2 - m_{13}^2 - m_{23}^2) \\ &\quad + \lambda^2 (J_1^2 J_2^2 + J_1^2 J_3^2 + J_2^2 J_3^2 - m_{12}^2 J_3^2 - m_{13}^2 J_2^2 - m_{23}^2 J_1^2) - J_1^2 J_2^2 J_3^2 \\ &= \lambda^6 - \lambda^4 (\text{tr}(J^2) - \|\mathbf{m}\|_2) + \lambda^2 (C_{J,2} - H_2) - \det(J^2). \end{aligned} \quad (18)$$

$$\begin{aligned} C_{J,i,j} &= J_1^i J_2^j + J_1^i J_3^j + J_2^i J_3^j, \\ C_{J,i} &= C_{J,i,i} \\ C_J &= C_{J,1} \\ H_2 &= (J_1 + J_2)(J_1 + J_3)(J_2 + J_3)H - C_J \|\mathbf{m}\|_2. \end{aligned}$$

- Reduce to a cubic equation (compute the roots explicitly)



## Schematical procedure

- Compute eigenvalues/eigenvectors of  $H_{\text{sympI}}$ :

$$H_{\text{sympI}} \begin{bmatrix} Y_1 & Y_2 \end{bmatrix} = \begin{bmatrix} Y_1 & Y_2 \end{bmatrix} \begin{bmatrix} \Lambda_+ & \\ & \Lambda_- \end{bmatrix}, \quad \text{Re } \Lambda_+ \geq 0,$$

(the eigenvectors need not be orthogonal and may be complex).  $Y_1, Y_2 \in \mathbb{R}^{6 \times 3}$ ,  $\Lambda_{\pm} \in \mathbb{R}^{3 \times 3}$ .

- Orthogonalize the eigenvectors (by Gram-Schmidt or QR),

$$[Y_1, Y_2] = QR,$$

so that

$$H_{\text{sympI}}Q = QR\Lambda R^{-1}$$

is the complex Schur form.

- Reduce to a real Schur form by considering real/imaginary part (complex Givens rotation).
- Compute  $S = Q_{21}Q_{11}^{-1}$ ,  $X = (M/2 + S)J^{-1}$ .

- 
- We don't need all the eigenvectors, just  $Y_1$ . Don't need  $R$ .
  - Avoid complex arithmetic altogether.



## The numerical DMV algorithm

- $\mathbf{m}_0 \mapsto hM_0 = h\hat{\mathbf{m}}_0$ .
- Compute the eigenvalues of  $H_{\text{symp1}} = H(M_k)$  solving for  $P(\lambda) = 0$  as in (18).
- For  $t_k = t_0 + kh$ ,  $k = 0, 1, 2 \dots$ 
  - Compute the (real) eigenvectors corresponding to  $\Lambda_+$ 
    - \* Compute 3 'quadratic' eigenvectors (3 matrix factorizations, LU/QR, with pivoting). No need to compute explicitly  $L$  or  $Q$ .
    - \* Compute the 'dependent' eigenvectors.
  - Orthogonalize the eigenspace
    - \* By (modified) Gram–Schmidt or QR. Only the  $Q$  factor is needed.
  - Compute  $S = Q_{21}Q_{11}^{-1}$ ,  $\omega_k = (M/2 + S)J^{-1}$ 
    - \* Update  $M_{k+1} = \omega_k^\top M_k \omega_k$
- Rescale  $\mathbf{m}_N \leftarrow M_N/h$ .

This algorithm produces an explicit method that is about 20 – 22 times more expensive than LP2, the explicit method of McLachlan and Reich.





# Concluding remarks

- **Explicit** algorithms to solve for the  $N = 3$  free rigid body
- The methods are up to 6th order, completely integrable, possible to increase to arbitrary order
- The cost of the method is about 10 – 22 times more expensive than the explicit LP2. The cheaper versions seem to be less stable especially for large step-size and long time computations
- Reconstruction equations: Find the configuration

$$X_{k+1} = X_k \omega_k^\top.$$

The complexive order is still 2 but the error is generally smaller. Backward error analysis reveals that some error terms cancel, however the components are reparametrized with 3 different time scales.

To obtain higher order reconstructions, one can use interpolation of the vector field  $X\Omega$  and its higher derivatives using the computed values of  $\Omega_k$ .

- Optimal step-size?





# References

- Cardoso, J. R. & Leite, F. S. (2003), 'The Moser–Veselov equation', *Lin. Alg. Applic.*
- Golub, G. H. & van Loan, C. F. (1989), *Matrix Computations*, 2nd edn, John Hopkins, Baltimore.
- Hairer, E., Lubich, C. & Wanner, G. (2002), *Geometric Numerical Integration*, number 31 in 'Springer Series in Computational Mathematics', Springer, Berlin.
- Marsden, J. E., Pekarsky, S. & Shkoller, S. (1999), 'Discrete Euler–Poincaré and Lie–Poisson equations', *Nonlinearity* **12**, 1647–1662.
- McLachlan, R. I. (1993), 'Explicit Lie–Poisson integration and the Euler equations', *Physical Review Letters* **71**, 3043–3046.
- Moser, J. & Veselov, A. P. (1991), 'Discrete Version of Some Classical Integrable Systems and Factorization of Matrix Polynomials', *Commun. Math. Phys.* **139**, 217–243.
- Munthe-Kaas, H. Z. & Krogstad, S. (2002), On enumeration problems in Lie–Butcher theory, Report in informatics, Institutt for informatikk, University of Bergen, Norway. To appear in *Future Generation Computer Systems*, Special issue.

